Modeling Blood Lead

John P. Christopher, Ph.D., D.A.B.T.
Staff Toxicologist
Human & Ecological Risk Division

30 September 2005

Modeling Blood Lead

- Construction of models
- DTSC's LeadSpread
- USEPA's Integrated Exposure, Uptake, and Biokinetc Model (IEUBK)
- Future Developments

For all chemicals except lead, safe levels in soil or waste correspond to <u>doses</u> which are not harmful. Safe doses of chemicals usually account for just one route of exposure.

- *e.g.* **oral RfD, inhalation RfC**

For lead, we choose a level in soil which results in 99% of a population having <u>blood lead levels</u> not exceeding 10 µg/dL, including all routes of exposure.

INPUTS TO BLOOD LEAD



Estimating Blood Lead

- **Both USEPA and DTSC use “slope factors” to estimate blood lead from an environment input.**
- **A slope factor is the change in blood lead for each µg/day of lead intake.**

Example of a Slope Factor

- Pocock *et al.* (1983) showed a linear increase in blood lead in thousands of men in the U.K. exposed to various levels of lead in drinking water.
- Blood lead in adult males increased 0.622 µg/dL for each additional 1 µg lead/day taken in from drinking water.
- If drinking water contains 5 µg/L,then the contribution to blood lead from drinking water is 3.1 µg/dL.

LeadSpread

- **Calculates cleanup levels for lead in in soil, given inputs for lead in air, water, and food.**
- **Calculates for 2-3 year old child or an adult.**
- **Inputs are central tendency values, not worst cases.**

INPUTS TO BLOOD LEAD



LeadSpread

The central tendency of blood lead is estimated:

$$\Sigma\ [Pb_{AIR}*SF_{AIR} + Pb_{WATER}*SF_{WATER} + Pb_{FOOD}*SF_{FOOD} + Pb_{SOIL}*SF_{SOIL} + Pb_{DUST}*SF_{DUST}]$$

LeadSpread

- **Blood lead in human populations has been shown repeatedly to be lognormally distributed. Lognormal distributions are not symmetrical, but the logs of the individual values are normally distributed.**



Lognormal Distirubtion



Normal Distirubtion

LeadSpread

- **Using the geometric mean (mean of log-transformed blood lead values) and an assumed geometric standard deviation (s.d of logs), any percentile of the lognormal distribution can be estimated:**

$$95^{th}\ \%ile = 10^{[\log(GM)+1.65*\log(GSD)]}$$

$$99^{th}\ \%ile = 10^{[\log(GM)+2.33*\log(GSD)]}$$

LEAD RISK ASSESSMENT SPREADSHEET

CALIFORNIA DEPARTMENT OF TOXIC SUBSTANCES CONTROL

USER'S GUIDE **to version 7**

INPUT	
MEDIUM	LEVEL
Lead in Air (µg/m^3)	0.028
Lead in Soil/Dust (µg/g)	146.0
Lead in Water (µg/L)	15
% Home-grown Produce	7%
Respirable Dust (µg/m^3)	1.5

OUTPUT							
	Percentile Estimate of Blood Pb (µg/dL)					PRG-99	PRG-95
	50th	90th	95th	98th	99th	(µg/g)	(µg/g)
BLOOD Pb, ADULT	1.5	2.8	3.3	4.0	4.6	694	1,080
BLOOD Pb, CHILD	3.3	6.1	7.2	8.8	10.0	146	247
BLOOD Pb, PICA CHILD	4.4	8.0	9.5	11.5	13.1	94	159
BLOOD Pb, OCCUPATIONAL	1.1	2.1	2.5	3.0	3.4	3,565	5,554

EXPOSURE PARAMETERS			
	units	adults	children
Days per week	days/wk	7	
Days per week, occupational		5	
Geometric Standard Deviation		1.6	
Blood lead level of concern (µg/dL)		10	
Skin area, residential	cm^2	5,700	2,900
Skin area occupational	cm^2	2,900	
Soil adherence	µg/cm^2-day	70	200
Dermal uptake constant	(µg/dL)/(µg/day)	9.6 E-5	
Soil ingestion	mg/day	50	100
Soil ingestion, pica	mg/day		200
Ingestion constant	(µg/dL)/(µg/day)	0.04	0.16
Bioavailability	unitless	0.44	
Breathing rate	m^3/day	20	6.8
Inhalation constant	(µg/dL)/(µg/day)	0.08	0.192
Water ingestion	L/day	1.3	0.4
Food ingestion	kg food/day	1.9	1.1
Lead in market basket	µg Pb/kg food	3.1	
Lead in home-grown produce	µg Pb/kg food	65.7	

Click here for REFERENCES

PATHWAYS						
ADULTS	Residential			Occupational		
	Pathway Contribution			Pathway Contribution		
Pathway	PEF	µg/dL	Percent	PEF	µg/dL	Percent
Soil Contact	3.8E-5	0.01	0%	1.4E-5	0.00	0%
Soil Ingestion	8.8E-4	0.13	8%	6.3E-4	0.09	8%
Inhalation, bkgrnd		0.05	3%		0.03	3%
Inhalation	2.5E-6	0.00	0%	1.8E-6	0.00	0%
Water Ingestion		0.78	51%		0.78	68%
Food Ingestion, bkgrnd		0.22	14%		0.24	21%
Food Ingestion	2.4E-3	0.35	23%			0%

CHILDREN	Typical Child			Child with Pica		
	Pathway Contribution			Pathway Contribution		
Pathway	PEF	µg/dL	Percent	PEF	µg/dL	Percent
Soil Contact	5.6E-5	0.01	0%		0.01	0%
Soil Ingestion	7.0E-3	1.03	31%	1.4E-2	2.06	47%
Inhalation	2.0E-6	0.00	0%		0.00	0%
Inhalation, bkgrnd		0.04	1%		0.04	1%
Water Ingestion		0.96	29%		0.96	22%
Food Ingestion, bkgrnd		0.51	15%		0.51	12%
Food Ingestion	5.5E-3	0.81	24%		0.81	18%

IEUBK

- **Models blood lead for children 6 months to 7 years old.**
- **IEUBK is an <u>uptake</u> model. It models blood lead from the <u>absorbed</u> dose.**
- **LeadSpread uses <u>applied</u> dose.**

IEUBK Similarities with LeadSpread

- **Both predict central tendency of blood lead, then generate a lognormal distribution using an assumed geometric standard deviation**
- **Both allow site-specific inputs.**

IEUBK Is More Detailed

- **Corrects for maternal contribution to blood lead.**
- **Corrects for non-linearities in absorption, distribution, and excretion of lead. [Slope factors allow only linear response.]**
- **Corrects for changing diet and water consumption with age.**

IEUBK Is More Detailed

- **Makes many pharmacokinetic corrections as children grow:**

  - increasing body size

  - decreasing G-I absorption of lead

  - storage of lead in bone

  - increasing bone mass

  - multiple types of bone

  - storage of lead in soft tissues

IEUBK Is More Detailed

- **Distinguishes between exposures to outside soil and inside dust.**
- **Allows user to build an extra pathway for special exposures, such as paint, cosmetics, or cookware.**
- **Allows exposure at more than one site.**

IEUBK Is More Detailed

- **Output includes the probability of a child exceeding a blood lead value of 10 µg/dL, given the specific inputs.**

LEAD MODEL FOR WINDOWS Version 1.0 Build 251

==========

Model Version: 1.0 Build 251
User Name:
Date:
Site Name:
Operable Unit:
Run Mode: Research

==========

The time step used in this model run: 1 - Every 4 Hours (6 times a day).

****** Air ******

Indoor Air Pb Concentration: 30.000 percent of outdoor.
Other Air Parameters:

Age	Time Outdoors (hours)	Ventilation Rate (m^3/day)	Lung Absorption (%)	Outdoor Air Pb Conc (ug Pb/m^3)
.5-1	1.000	2.000	32.000	0.100
1-2	2.000	3.000	32.000	0.100
2-3	3.000	5.000	32.000	0.100
3-4	4.000	5.000	32.000	0.100
4-5	4.000	5.000	32.000	0.100
5-6	4.000	7.000	32.000	0.100
6-7	4.000	7.000	32.000	0.100

****** Diet ******

Age	Diet Intake (ug/day)
.5-1	5.530
1-2	5.780
2-3	6.490
3-4	6.240
4-5	6.010
5-6	6.340
6-7	7.000

****** Drinking Water ******

Water Consumption:

Age	Water (L/day)
.5-1	0.200
1-2	0.500
2-3	0.520
3-4	0.530
4-5	0.550
5-6	0.580
6-7	0.590

Drinking Water Concentration: 4.000 ug Pb/L

****** Soil & Dust ******

Multiple Source Analysis Used
Average multiple source concentration: 150.000 ug/g

Mass fraction of outdoor soil to indoor dust conversion factor: 0.700
Outdoor airborne lead to indoor household dust lead concentration: 100.000
Use alternate indoor dust Pb sources? No

Age	Soil (ug Pb/g)	House Dust (ug Pb/g)
.5-1	200.000	150.000
1-2	200.000	150.000
2-3	200.000	150.000
3-4	200.000	150.000
4-5	200.000	150.000
5-6	200.000	150.000
6-7	200.000	150.000

****** Alternate Intake ******

Age	Alternate (ug Pb/day)
.5-1	0.000
1-2	0.000
2-3	0.000
3-4	0.000
4-5	0.000
5-6	0.000
6-7	0.000

****** Maternal Contribution: Infant Model ******

Maternal Blood Concentration: 2.500 ug Pb/dL

****
CALCULATED BLOOD LEAD AND LEAD UPTAKES:
****

Year	Air (ug/dL)	Diet (ug/day)	Alternate (ug/day)	Water (ug/day)
.5-1	0.021	2.553	0.000	0.369
1-2	0.034	2.647	0.000	0.916
2-3	0.062	3.002	0.000	0.962
3-4	0.067	2.919	0.000	0.992
4-5	0.067	2.863	0.000	1.048
5-6	0.093	3.040	0.000	1.112
6-7	0.093	3.366	0.000	1.135

Year	Soil+Dust (ug/day)	Total (ug/day)	Blood (ug/dL)
.5-1	4.061	7.004	3.8
1-2	6.399	9.997	4.2
2-3	6.462	10.488	3.9
3-4	6.536	10.514	3.7
4-5	4.930	8.908	3.1
5-6	4.467	8.712	2.7
6-7	4.231	8.826	2.5

IEUBK Example

	Input Parameters					Output from Model	
	Lead in Soil (mg/kg)	Lead in Dust (mg/kg)	Lead in Water (µg/L)	Age (months)	Other Inputs	Geometric Mean Blood Lead (µg/dL)	% Above 10 µg/dL
Run 1	Default	Default	Default	0-84	Default	3.4	1.1
Run 2	500	500	Default	0-84	Default	6.6	18.8
Run 3	500	500	50	0-84	Default	9.4	44.34
Run 4	1,000	1,000	Default	0-84	Default	10.7	55.5
Run 5	1,000	1,000	50	0-84	Default	13	71.1

Uncertainties

- **The largest uncertainty in either model is the assumed estimate of variance, the geometric standard deviation.**
- **The default value of 1.6 is actually a combination of variability in:**
  - **- concentration in soil**
  - **- exposure parameters**
  - **- pharmacokinetcs**

Uncertainties

- **We would like our estimate of variability to be confined to “inter-individual variability”, which does not include the variability in concentration in soil.**
- **Separating out the three types of variability is not a simple task.**
- **The default value of 1.6 errs on the high side, which is the health-protective direction.**

Conclusions

- IEUBK provides the most detailed modeling available for blood lead in children.
- LeadSpread provides a first approximation of IEUBK results for the 2-3 year old child.